

# NEW JERSEY ADULT MOSQUITO SURVEILLANCE

Report for 11 August to 17 August 2013, CDC Week 33

Prepared by Lisa M. Reed, Scott Crans and Mark Robson  
Center for Vector Biology



This New Jersey Agricultural Experiment Station report is supported by Rutgers University, Hatch funds, funding from the NJ State Mosquito Control Commission and with the participation of the 21 county mosquito control agencies of New Jersey.


**Summary Table – Week 33**

Region	<i>Aedes vexans</i>			<i>Culex Mix</i>			<i>Coquillettidia perturbans</i>			<i>Aedes sollicitans</i>		
	This Week	Average*	Increase	This Week	Average*	Increase	This Week	Average*	Increase	This Week	Average*	Increase
Agricultural	<b>0.31</b>	<b>1.84</b>	0	<b>0.69</b>	<b>2.34</b>	0	<b>0.02</b>	<b>0.20</b>	0	<b>0.00</b>	<b>0.15</b>	0
Coastal	<b>0.30</b>	<b>2.38</b>	0	<b>3.39</b>	<b>6.02</b>	0	<b>0.04</b>	<b>0.16</b>	0	<b>0.54</b>	<b>8.37</b>	0
Delaware Bayshore	<b>0.23</b>	<b>1.31</b>	0	<b>2.91</b>	<b>9.99</b>	0	<b>0.03</b>	<b>0.40</b>	0	<b>0.00</b>	<b>5.08</b>	0
Delaware River Basin	nd	<b>4.42</b>	0	nd	<b>2.34</b>	0	nd	<b>0.69</b>	0	nd	<b>0.00</b>	0
New York Metro	<b>0.17</b>	<b>5.03</b>	0	<b>2.13</b>	<b>8.24</b>	0	<b>0.00</b>	<b>0.12</b>	0	<b>0.17</b>	<b>0.11</b>	2
North Central Rural	<b>1.27</b>	<b>1.04</b>	1	<b>1.02</b>	<b>1.14</b>	0	<b>0.00</b>	<b>0.01</b>	0	<b>0.00</b>	<b>0.00</b>	0
Northwest Rural	<b>0.97</b>	<b>9.66</b>	0	<b>5.17</b>	<b>5.65</b>	0	<b>0.05</b>	<b>0.70</b>	0	<b>0.00</b>	<b>0.00</b>	0
Philadelphia Metro	nd	<b>8.48</b>	0	nd	<b>2.77</b>	0	nd	<b>0.05</b>	0	nd	<b>0.00</b>	0
Pinelands	<b>0.14</b>	<b>1.10</b>	0	<b>1.30</b>	<b>3.30</b>	0	<b>0.12</b>	<b>0.35</b>	0	<b>0.00</b>	<b>0.12</b>	0
Suburban Corridor	<b>0.47</b>	<b>8.40</b>	0	<b>0.75</b>	<b>1.43</b>	0	<b>0.03</b>	<b>0.56</b>	0	<b>0.00</b>	<b>0.02</b>	0

\*Averages represent data from, at most, the previous 5 years. Increase is a scale of current values from historical values where no difference or a decrease is represented by 0 (blue), up to 50% greater difference by 1 (green), up to 100% greater difference by 2 (yellow), up to 150% greater difference by 3 (orange) and greater than 150% increase by 4 (red). White cells in the increase column denote increases from an historic zero and thus no value can be appropriately given. nd=no data reported.


State Summary: Mosquito populations were only moderately above in two regions (*Aedes vexans* in the North Central Rural and *Aedes sollicitans* in the New York Metropolitan region), but this may change as data continues to come in.

## Climate Factors




The three figures show the interpolation of average maximum ( $^{\circ}\text{F}$ ) and minimum temperature ( $^{\circ}\text{F}$ ) and total precipitation (inches) from 1 August to 15 August, 2013 in New Jersey. Data points are from about 43 weather stations maintained through the New Jersey Weather & Climate Network and the State Climatologist. Interpolation between points was performed using ArcMap 10.1.

**The Species Graphs:** The species graph pages include a graph with two plots for each of the ten regions defined on the first page (Agricultural, Coastal, Delaware Bayshore, Delaware River, New York Metro, North-Central, Northwestern, Philadelphia Metro, Pinelands, and Suburban Corridor). Below is an example of one graph from one species within one region. The bar plot show the average number of mosquitoes per trap within the region (weekly means) and line plots show the historical trend as the average number of mosquitoes from the previous 5 years (5-year average). In general, historical data are running means from the previous 5 years, but on occasion, will include data from fewer years. Adjustments are made to account for year discrepancies. Data for this week are from Cape May, Essex, Monmouth, Morris, Ocean, Sussex, Union and Warren counties. Data for the previous week(s) are from Atlantic, Bergen, Burlington, Camden, Cape May, Essex, Hudson, Hunterdon, Monmouth, Morris, Ocean, Somerset, Sussex, Union and Warren counties.




# Aedes vexans - Fresh Floodwater Species

## Multivoltine Aedine (Ae. vexans Type)




# Culex Mix – Permanent Water Species

## Multivoltine Culex/Anopheles (*Cx. pipiens* Type)




# *Culiseta melanura* – Miscellaneous Group

## Unique (*Cs. melanura* Type)




# *Aedes sollicitans* - Salt Floodwater Species

## Multivoltine Aedine (*Ae. sollicitans* Type)



# *Coquillettidia perturbans*

## Monotypic (*Coquillettidia perturbans* Type)




WNV

EEE


**Top Ten Mosquito Species/Region -** ■ Ae. albopictus, ■ Ae. japonicus (invasives); ■ Cs. melanura or Cx. erraticus ■ Coq. perturbans

Note: In early season when fewer species are caught, graphs may show less than ten species/region or 25 statewide.




## Agricultural



Total # mosquitoes



## Coastal


Total # mosquitoes






## New York Metropolitan

Total # mosquitoes




## North Central Rural

Total # mosquitoes




## Northwest Rural

Total # mosquitoes




## Philadelphia Metropolitan

Total # mosquitoes



## Pinelands

Total # mosquitoes



## Suburban Corridor

Total # mosquitoes

