

# NEW JERSEY ADULT MOSQUITO SURVEILLANCE

Report for 7 June to 13 June 2015, CDC Week 23

Prepared by Lisa M. Reed, Scott Crans and Mark Robson

Center for Vector Biology



This New Jersey Agricultural Experiment Station report is supported by Rutgers University, Hatch funds, funding from the NJ State Mosquito Control Commission and with the participation of the 21 county mosquito control agencies of New Jersey.

**Summary Table – Week 23**


Region	<i>Aedes vexans</i>			<i>Culex Mix</i>			<i>Coquillettidia perturbans</i>			<i>Aedes sollicitans</i>		
	This Week	Average*	Increase	This Week	Average*	Increase	This Week	Average*	Increase	This Week	Average*	Increase
Agricultural	0.23	2.49	0	3.33	1.88	2	0.33	0.49	0	0.10	0.02	4
Coastal	0.00	2.27	0	0.52	3.53	0	0.00	0.28	0	0.00	3.11	0
Delaware Bayshore	0.34	2.78	0	2.86	12.54	0	1.91	5.27	0	0.49	1.22	0
Delaware River Basin	0.43	1.79	0	6.21	0.90	4	0.36	0.16	3	0.00	0.00	0
New York Metro	0.07	1.77	0	1.06	1.34	0	0.03	0.22	0	0.00	0.02	0
North Central Rural	0.04	0.25	0	0.71	0.75	0	0.00	0.03	0	0.00	0.00	0
Northwest Rural	0.03	5.49	0	0.00	2.18	0	0.00	0.44	0	0.00	0.00	0
Philadelphia Metro	nd	3.92	0	nd	5.14	0	nd	0.80	0	nd	0.00	0
Pinelands	0.04	0.82	0	0.06	1.48	0	0.00	0.52	0	0.00	0.03	0
Suburban Corridor	0.12	5.04	0	0.56	1.83	0	0.00	0.48	0	0.00	0.01	0

\*Averages represent data from, at most, the previous 5 years. Increase is a scale of current values from historical values where no difference or a decrease is represented by 0 (blue), up to 50% greater difference by 1 (green), up to 100% greater difference by 2 (yellow), up to 150% greater difference by 3 (orange) and greater than 150% increase by 4 (red). White cells in the increase column denote increases from an historic zero and thus no value can be appropriately given. nd=no data reported.

**State Summary:** *Culex* species (likely *Cx. restuans*) in the Agricultural region and especially the Delaware River Basin are above historical averages. *Coquillettidia perturbans* populations are also elevated in the Delaware River. *Ae. sollicitans* numbers, although low, are also above historical values in the Agricultural region.

**Note:** CDC weeks corrected from Excel week formula from previous reports.


## Climate Factors



The three figures show the interpolation of average maximum ( $^{\circ}\text{F}$ ) and minimum temperature ( $^{\circ}\text{F}$ ) and total precipitation (inches) for 30 days prior to 12 June 2015 in New Jersey. Data points are from about 45 weather stations maintained through the New Jersey Weather & Climate Network and the State Climatologist. Interpolation between points was performed using ArcMap 10.1.

**The Species Graphs:** The species graph pages include a graph with two plots for each of the ten regions defined on the first page (Agricultural, Coastal, Delaware Bayshore, Delaware River, New York Metro, North-Central, Northwestern, Philadelphia Metro, Pinelands, and Suburban Corridor). Below is an example of one graph from one species within one region. The bar plot show the average number of mosquitoes per trap within the region (weekly means) and line plots show the historical trend as the average number of mosquitoes from the previous 5 years (5-year average). In general, historical data are running means from the previous 5 years, but on occasion, will include data from fewer years. Adjustments are made to account for year discrepancies. Data for this week are from Cumberland, Hudson, Hunterdon, Monmouth, Morris, Salem, Union and Warren counties. Data for the previous week are from Atlantic, Bergen, Cape May, Cumberland, Hudson, Hunterdon, Mercer, Monmouth, Morris, Ocean, Salem, Sussex, Union and Warren counties.

## Weekly Means Against 5-year Average




# Aedes vexans - Fresh Floodwater Species Multivoltine Aedine (Ae. vexans Type)


<p><b>Agricultural</b></p>	<p><b>Coastal</b></p>	<p><b>Delaware Bayshore</b></p>	<p><b>Delaware River Basin</b></p>
<p><b>New York Metro</b></p>	<p><b>North Central Rural</b></p>	<p><b>Northwestern Rural</b></p>	<p><b>Philadelphia Metro</b></p>
<p><b>Pinelands</b></p>	<p><b>Suburban Corridor</b></p>	<p><b>Comments</b></p> <p><i>Aedes vexans</i> populations continue to remain below historical averages for most regions. Despite significant rainfall, abnormally dry and moderate drought conditions continue to exist for the northern half of the state. Water runoff may have been too fast for adequate ground absorption. But current warm weather may quicken any larval development in any remaining habitat and an increase in adults could occur in the coming week.</p>	

# Culex Mix – Permanent Water Species


## Multivoltine *Culex/Anopheles* (*Cx. pipiens* Type)



## *Culiseta melanura* – Miscellaneous Group Unique (*Cs. melanura* Type)




## Aedes sollicitans - Salt Floodwater Species Multivoltine Aedine (Ae. sollicitans Type)



# Coquillettidia perturbans

## Monotypic (*Coquillettidia perturbans* Type)






WNV

EEE


**Top Ten Mosquito Species/Region** - ■ *Ae. albopictus*, ■ *Ae. japonicus* (invasives); ■ *Cs. melanura* or *Cx. erraticus* ■ *Coq. perturbans*

Note: In early season when fewer species are caught, graphs may show less than ten species/region or 25 statewide.




## Agricultural



Total # mosquitoes



## Coastal


Total # mosquitoes






## New York Metropolitan

Total # mosquitoes




## North Central Rural

Total # mosquitoes




## Northwest Rural

Total # mosquitoes




## Philadelphia Metropolitan

Total # mosquitoes



## Pinelands

Total # mosquitoes



## Suburban Corridor

Total # mosquitoes

