

NEW JERSEY ADULT MOSQUITO SURVEILLANCE Report

July 19 to July 25, CDC Week 30

Prepared by Lisa M. Reed and Dina Fonseca
Center for Vector Biology



This New Jersey Agricultural Experiment Station report is supported by Rutgers University, Hatch funds, funding from the NJ State Mosquito Control Commission and with the participation of the 21 county mosquito control agencies of New Jersey.


Summary Table – Week 30

Region	<i>Aedes vexans</i>			<i>Culex Mix</i>			<i>Coquillettidia perturbans</i>			<i>Aedes sollicitans</i>		
	This Week	Average*	Increase	This Week	Average*	Increase	This Week	Average*	Increase	This Week	Average*	Increase
Agricultural	12.12	3.60	4	8.29	6.98	1	0.21	0.56	0	10.76	2.45	4
Coastal	3.16	2.38	1	1.70	7.29	0	0.00	0.29	0	3.08	1.51	3
Delaware Bayshore	0.95	1.05	0	9.07	12.36	0	0.00	1.09	0	14.97	2.25	4
Delaware River Basin	45.43	9.91	4	9.64	4.59	3	7.21	0.60	4	0.00	0.01	0
New York Metro	0.63	2.78	0	1.84	6.52	0	0.06	0.07	0	0.01	0.60	0
North Central Rural	0.04	0.21	0	0.02	0.48	0	0.14	0.26	0	0.00	0.00	0
Northwest Rural	0.06	11.02	0	0.09	4.71	0	0.00	0.53	0	0.00	0.00	0
Philadelphia Metro	0.00	4.33	0	0.00	3.25	0	0.00	0.03	0	0.00	0.00	0
Pinelands	1.84	1.39	1	2.00	3.22	0	0.58	1.33	0	0.30	0.09	4
Suburban Corridor	0.11	1.73	0	0.16	0.97	0	0.00	0.19	0	0.00	0.02	0

*Averages represent data from, at most, the previous 5 years. Increase is a scale of current values from historical values where no difference or a decrease is represented by 0 (blue), up to 50% greater difference by 1 (green), up to 100% greater difference by 2 (yellow), up to 150% greater difference by 3 (orange) and greater than 150% increase by 4 (red). White cells in the increase column denote increases from an historic zero and thus no value can be appropriately given. nd=no data reported.

State Summary: This week's data represents 10 out of 21 counties. Significant populations for all 4 pestiferous species are seen, including activity from floodwater species. *Aedes vexans* populations were significantly higher than the 5 year running average in the Agricultural and Delaware River Basin regions and mildly elevated in the Coastal and Pinelands regions. Similarly, *Aedes sollicitans* were also significantly elevated in the Agricultural and Delaware River Basin regions as well as in the Pinelands and elevated in the Coastal region. *Culex Mix* was elevated in the Delaware River Basin region and mildly above average in the Agricultural region. *Coquillettidia perturbans* were significantly higher in the Delaware River Basin region.


Climate Factors




The three figures show the interpolation of average maximum (°F) and minimum temperature (°F) and total precipitation (inches) for 30 days prior to 24 July 2020 in New Jersey. Data points are from about 50 weather stations maintained through the New Jersey Weather & Climate Network and the State Climatologist. Interpolation between points was performed using ArcMap 10.1.

The Species Graphs: The species graph pages include a graph with two plots for each of the ten regions defined on the first page (Agricultural, Coastal, Delaware Bayshore, Delaware River, New York Metro, North-Central, Northwestern, Philadelphia Metro, Pinelands, and Suburban Corridor). Below is an example of one graph from one species within one region. The bar plot show the average number of mosquitoes per trap within the region (weekly means) and line plots show the historical trend as the average number of mosquitoes from the previous 5 years (5-year average). In general, historical data are running means from the previous 5 years, but on occasion, will include data from fewer years. Adjustments are made to account for year discrepancies. Data for this week are from Atlantic, Cumberland, Hudson, Hunterdon, Monmouth, Morris, Ocean, Salem, Somerset, and Warren counties. Data for the previous week are from Atlantic, Cape May, Cumberland, Hudson, Hunterdon, Monmouth, Morris, Ocean, Salem, Somerset, and Warren counties.


Weekly Means Against 5-year Average




Aedes vexans - Fresh Floodwater Species Multivoltine Aedine (Ae. vexans Type)




Culex Mix – Permanent Water Species Multivoltine *Culex/Anopheles* (*Cx. pipiens* Type)



Culiseta melanura – Miscellaneous Group Unique (*Cs. melanura* Type)




Aedes sollicitans - Salt Floodwater Species Multivoltine Aedine (Ae. sollicitans Type)



Coquillettidia perturbans

Monotypic (*Coquillettidia perturbans* Type)




WNV

EEE


Top Ten Mosquito Species/Region - ■ *Ae. albopictus*, ■ *Ae. japonicus* (invasives); ■ *Cs. melanura* or *Cx. erraticus* ■ *Coq. perturbans*

Note: In early season when fewer species are caught, graphs may show less than ten species/region or 25 statewide.




Agricultural

Total # mosquitoes




Coastal

Total # mosquitoes




Delaware Bayshore

Total # mosquitoes




Delaware River Basin

Total # mosquitoes




New York Metropolitan

Total # mosquitoes




North Central Rural

Total # mosquitoes




Northwest Rural

Total # mosquitoes




Philadelphia Metropolitan

Total # mosquitoes



Pinelands

Total # mosquitoes



Suburban Corridor

Total # mosquitoes

