

# VECTOR SURVEILLANCE IN NEW JERSEY

## EEE, WNV, SLE and LAC

Prepared by Lisa M. Reed, Scott Crans and Mark Robson

Center for Vector Biology, Rutgers University

CDC WEEK 23: Beginning May to 4 June, 2014

Data Downloaded 11:58 pm 4 June 2014



This New Jersey Agricultural Experiment Station report is supported by Rutgers University, Hatch funds, funding from the NJ State Mosquito Control Commission and with the participation of the Department of Health, Department of Agriculture and of the 21 county mosquito control agencies of New Jersey.

### *Culiseta melanura* and Eastern Equine Encephalitis

SITE/Boxes	Inland or Coastal	Historic Population Mean	Current Weekly Mean	Total Tested* (Collected)	Total Pools Tested* (Submitted)	EEE Isolation Pools	MFIR
Bass River (Burlington Co.)/5	Coastal	0.10	0.00	1			
Green Bank (Burlington Co.)/25	Coastal	0.91	0.16	23			
Corbin City (Atlantic Co.)/25	Coastal	1.10	0.64*	16			
Dennisville (Cape May Co.)/50	Coastal	3.87	0.22	11			
Winslow (Camden Co.)/50	Inland	2.13	3.50	140			
Centerton (Salem Co.)/50	Inland	2.07	0.86	43			
Turkey Swamp (Monmouth Co.)/44	Inland	0.52	0.04	14			
Glassboro (Gloucester Co.)/50	Inland	0.52	0.42	21			

\*Current week (in parentheses) results pending. Corbin City from Week 22.

**Remarks:** The resting boxes at the traditional sites have been placed out in the beginning of May to condition and collections began the last week of May. Currently, sites are active with mosquitoes using the resting boxes (see population graphs page 3).

**Traditional Resting Box Sites:** 269 *Culiseta melanura* from the resting boxes are currently in the system and are being updated to the state site. These pending results will be report next week.

**Additional *Cs. melanura* trapped by counties**

\*traps with positives indicated in **BOLD**.

County	Trap types*	Number collected (pools)	Number of positive pools	MFIR
<b>TOTAL</b>				

**Additional *Cs. melanura*:** Counties submit additional pools of *Cs. melanura* caught in other trap types as well as resting boxes. These results will be reported when available.

Species other than <i>Cs. melanura</i>	Pools	Mosquitoes	Positives	MFIR
<i>Aedes albopictus</i>				
<i>Aedes atlanticus</i>				
<i>Aedes canadensis canadensis</i>				
<i>Aedes cantator</i>				
<i>Aedes infirmatus</i>				
<i>Aedes japonicus</i>				
<i>Aedes sollicitans</i>				
<i>Aedes sticticus</i>				
<i>Aedes taeniorhynchus</i>				
<i>Aedes triseriatus</i>				
<i>Aedes vexans</i>				
<i>Anopheles bradleyi</i>				
<i>Anopheles crucians</i>				
<i>Anopheles punctipennis</i>				
<i>Anopheles quadrimaculatus</i>				
<i>Coquillettidia perturbans</i>				
<i>Culex erraticus</i>				
<i>Culex pipiens</i>				
<i>Culex restuans</i>				
<i>Culex salinarius</i>				
<i>Culex</i> spp.				
<i>Psorophora columbiae</i>				
State Total				

**Additional Species:** Counties submit additional pools of species other than *Cs. melanura* for EEE virus testing. Last year, in addition to *Cs. melanura*, other species that tested positive included *Culex erraticus*, *Cx. pipiens*, *Cx. salinarius*, *Coquillettidia perturbans* and *Anopheles bradleyi*. These additional species will be reported when available.


**Horses and Humans:** Currently there is no reported horse or human cases. Last year, a total of 5 horses developed EEE, resulting in euthanasia. They were either unvaccinated or with unknown/incomplete vaccination histories.

There were no reported human cases last year.

The graph to the right shows the end of year activity for all mosquitoes tested positive for EEE in 2013.

**Horses and Vaccinations:** The fate of unvaccinated equids reinforces the necessity of maintaining a vaccination schedule for arboviruses. For vaccination schedules recommended by the American Association of Equine Practices, see: [http://www.aaep.org/vaccination\\_guidelines.htm](http://www.aaep.org/vaccination_guidelines.htm)


Counties with all mosquito EEE activity




# Culiseta melanura Population Graphs

## Coastal


BASS RIVER (Burlington Co.)




GREEN BANK (Burlington Co.)



CORBIN CITY (Atlantic Co.)




DENNISVILLE (Cape May Co.)




## Inland


WINSLOW (Camden Co.)




CENTERTON (Salem Co.)




TURKEY SWAMP (Monmouth Co.)



GLASSBORO (Gloucester Co.)



The traditional resting box sites monitoring indicate *Cs. melanura* activity at all sites. Numbers appear high at the Winslow site but are within the standard deviation for the average activity over the past five years. The same can be said for the rest of the sites with regard to the seemingly low numbers. Populations are also currently being compiled from light trap data for the adult mosquito surveillance reports that can sometimes clarify the population status at different sites.

 = Positive pool(s) detected (red = melanura, purple = other).

EEE in US (2014 cumulative cases): (Black or Red = previous + new reported cases occurring)

- equine: 7(FL)
- mosquito pools:
- sentinel: 21(FL)
- human:

### West Nile Virus Positive Organisms in US

West Nile in US (2014 cumulative cases): Single black values indicate no change from previous week. Black values / red values equals previous week/**New totals**.  
 Note: Data reported by all states should be considered provisional and subject to change. Sources for this table can be found [here](#).

	Birds	Mosquito Pools	Sentinels	Horses	Humans
Alabama					
Alaska					
Arizona	1	1			
Arkansas					
California	65	23	1		
Colorado					
Connecticut					
Delaware					
DC					
Florida			4		
Georgia					
Hawaii					
Idaho					
Illinois	1	1			
Indiana					
Iowa					
Kansas					
Kentucky					
Louisiana					
Maine					
Maryland					
Mass.					
Michigan					
Minnesota					
Mississippi					2
Missouri					

	Birds	Mosquito Pools	Sentinels	Horses	Humans
Montana					
Nebraska					
Nevada					
New Hampshire					
New Jersey		1			
New Mexico					
New York					
North Carolina					
North Dakota					
Ohio					
Oklahoma					
Oregon					
Pennsylvania		5			
Rhode Island					
South Carolina					
South Dakota					
Tennessee					
Texas					
Utah					
Vermont					
Virginia					
Washington					
West Virginia					
Wisconsin	2				
Wyoming					

\* Can include other species (e.g., dogs, cows) reported positive.

Protocol: New Jersey Department of Health (NJDH Public Health Environmental and Agricultural Laboratories, PHEAL) and the Cape May County Department of Mosquito Control tests mosquito pools using RT-PCR Taqman techniques.

### Mosquito Species Submitted and Tested for West Nile Virus Testing through 4 June 2014

Species	Pools	Mosquitoes	Positives	MFIR
<i>Aedes canadensis canadensis</i>	1	4		
<i>Aedes japonicus</i>	4	7		
<i>Culex pipiens</i>	20	732		
<i>Culex spp.</i>	27	1019	1	0.981
<i>Culiseta melanura</i>	1	16		
<b>State Total</b>	<b>53</b>	<b>1778</b>	<b>1</b>	<b>0.562</b>

**Remarks:** To date, 53 pools of 1778 mosquitoes from 5 species have been tested, with 1 positive pool detected. First positive was detected in a Mixed *Culex* pool collected on 20 May in Camden County.


**Humans, Horses and Wild Birds:** To date, no human cases have been reported. For further information, see <http://www.state.nj.us/health/cd/westnile/techinfo.shtml>.

Last year the first (and only) horse was detected in early October. To date, no horses have been reported. As with EEE, most horses that develop West Nile have no or uncertain vaccination histories.

Bird testing began in mid-April. No positive birds have been reported. To date, 19 birds have been tested. Species includes: Fish Crow (*C. ossifragus* 0/5), Hawk/Raptor (0/2) and other avian species (0/12). Counties (**positives**) submitting birds are Bergen, Burlington, Essex, Mercer, Monmouth, Morris, Ocean, Salem and Warren.

### WNV Results by County through 4 June 2014

County	Species	Pools	Mosquitoes	Positives	MFIR
<b>Atlantic</b>		<b>2</b>	<b>20</b>		
	<i>Aedes canadensis canadensis</i>	1	4		
	<i>Culiseta melanura</i>	1	16		
<b>Camden</b>		<b>16</b>	<b>313</b>	<b>1</b>	<b>3.195</b>
	<i>Aedes japonicus</i>	4	7		
	<i>Culex spp.</i>	12	306	1	3.268
<b>Gloucester</b>		<b>20</b>	<b>732</b>		
	<i>Culex pipiens</i>	20	732		
<b>Hunterdon</b>		<b>15</b>	<b>713</b>		
	<i>Culex spp.</i>	15	713		
<b>Grand Total</b>		<b>53</b>	<b>1778</b>	<b>1</b>	<b>0.562</b>



Cumulative WNV activity in 2013.      WNV activity to 4 November 2014.      WNV activity last week, 2014.

### Saint Louis Encephalitis (SLE) to 4 June 2014.

New Jersey will be selectively testing for SLE this year. SLE has had previous activity in New Jersey, most notably in 1964 and 1975 (CDC's SLE [website](#)), the latter prompting the surveillance reporting by Rutgers. SLE is a flavivirus and has a similar transmission pattern to West Nile, with *Culex* species as the predominant vectors.

No pools have been reported for SLE in 2014.

County	Species	Pools	Mosquitoes	Positives	MFIR
<b>Grand Total</b>					

### La Crosse Encephalitis (LAC) through 4 June 2014.

New Jersey will be selectively testing for La Crosse (LAC) virus this year. New Jersey has had 3 cases of this encephalitic disease since 1964 (see CDC's LAC [website](#)). The mortality is low but like other encephalitides, LAC can have both personal (lasting neurological sequelae) and economic impacts. LAC is a bunyavirus with a transmission cycle involving mosquitoes such as *Aedes triseriatus* and small mammals such as squirrels and chipmunks. LAC can not only infect *Aedes albopictus* but transovarial transmission was also demonstrated. (Tesh and Gubler 1975 Laboratory studies of transovarial transmission of La Crosse and other arboviruses by *Aedes albopictus* and *Culex fatigans*. American Journal of Tropical Medicine and Hygiene 24(5):876-880).

No pools have been reported for LAC in 2014.

County	Species	Pools	Mosquitoes	Positives	MFIR
<b>Grand Total</b>					